# Author Index


Aarden, L.A., see Yap, S.H. (1091) 405
Ailhaud, G., see Catalioto, R.-M. (1091) 364
Al-Hassan, J.M., see Summers, B. (1091) 329
Albano, E., Bellomo, G., Parola, M., Carini, R. and Dianzani, M.U.
Stimulation of lipid peroxidation increases the intracellular calcium content of isolated hepatocytes (1091) 310
Alpers, D.H., see Eliakim, R. (1091) 1
Antoni, F., see Farkas, G. (1091) 81
Arai, H., see Kume, N. (1091) 63
Arnold, R., see Fehmann, H.-C. (1091) 356

Bächle, R., see Fehmann, H.-C. (1091) 356
Baker, E., see Richardson, D.R. (1091) 294
Banga, H.S., Halenda, S.P. and Feinstein, M.B.
Potentiation of arachidonic acid release by phorbol myristate acetate in platelets is not due to inhibition of arachidonic acid uptake or incorporation into phospholipids (1091) 115
Baxter, M.A., Bunce, C.M., Lord, J.M., French, P.J., Michell, R.H. and Brown, G.
Changes in inositol transport during DMSO-induced differentiation of HL60 cells towards neutrophils (1091) 158
Bellomo, G., see Albano, E. (1091) 310
Berton, G., Sorio, C., Laudanna, C., Menegazzi, M., Carcereri De Prati, A. and Suzuki, H.
Activation of human monocyte-derived macrophages by interferon γ is accompanied by increase polymerase activity of poly(ADP-ribose) (1091) 101
Bertrams, A. and Ziegler, K.
Hepatocellular uptake of peptides by bile acid transporters: relationship of carrier-mediated transport of linear peptides with renin-inhibiting activity to multispecific bile acid carriers (1091) 337
Best, L., see Lynch, A.M. (1091) 141
Bijzet, J., see Yap, S.H. (1091) 405
Blachier, F., Mourtada, A., Gomis, R., Sener, A. and Malaisse, W.J.
Metabolic and secretory response of parotid cells to cationic amino acids. Uptake and catabolism of L-arginine and L-ornithine (1091) 151
Bockman, R.S., see Rapuano, B.E. (1091) 374
Borin, G., see Brunati, A.M. (1091) 123
Bowmaster, T.A., Davis, C.C. and Krauthamer, V.
Excitation and detection of action potential-induced fluorescence changes through a single monomode optical fiber (1091) 9
Braquet, P., see Zor, U. (1091) 385
Brown, G., see Baxter, M.A. (1091) 158
Brunati, A.M., Donella-Deana, A., Ralph, S., Marchiori, F., Borin, G., Fischer, S. and Pinna, L.A.
Stimulation by NaCl, polylysine and heparin of two forms of spleen tyrosine protein kinase immunologically related with the protein expressed by *lyn* oncogene (1091) 123
Buday, L., see Farkas, G. (1091) 81
Bunce, C.M., see Baxter, M.A. (1091) 158
Burgess, B., see Panus, P.C. (1091) 303

Cambrey, A.D., see McAnulty, R.J. (1091) 231
Campa, J.S., see McAnulty, R.J. (1091) 231
Carcereri De Prati, A., see Berton, G. (1091) 101
Carini, R., see Albano, E. (1091) 310
Carrera, C.J., see Carson, D.A. (1091) 22
Carson, D.A., Carrera, C.J., Wasson, D.B. and Iizasa, T.
Deoxyadenosine-resistant human T lymphoblasts with elevated 5′-nucleotidase activity (1091) 22
Catalioto, R.-M., Gaillard, D., Maclouf, J., Ailhaud, G. and Negrel, R.
Autocrine control of adipose cell differentiation by prostacyclin and $PGF_{2\alpha}$ (1091) 364
Charbonneau, M., see Grandin, N. (1091) 242
Chun, A., see Summers, B. (1091) 329
Criddle, R.S., see Summers, B. (1091) 329

Davis, C.C., see Bowmaster, T.A. (1091) 9
Deana, A.D., see Leader, D.P. (1091) 426
DelValle, J., see Tsunoda, Y. (1091) 251
Deslauriers, C., Powell, A.D., Fuchs, K. and Pauls, K.P.
Flow cytometric characterization and sorting of cultured *Brassica napus* microspores (1091) 165
Dhar, A. and Shukla, S.D.
Release of a membrane surface glycoprotein from human platelets by phosphatidylinositol specific phospholipase(s) C (1091) 15
Dianzani, M.U., see Albano, E. (1091) 310
Dolphin, A.C.
Regulation of calcium channel activity by GTP binding proteins and second messengers (1091) 68
Donella-Deana, A., see Brunati, A.M. (1091) 123
Duhm, J., see Engelmann, B. (1091) 261

Eliakim, R., Mahmood, A. and Alpers, D.H.
Rat intestinal alkaline phosphatase secretion into lumen and serum is coordinately regulated (1091) 1
Elias, E., see Garner, C.M. (1091) 41
Engelmann, B., Schumacher, U. and Duhm, J.
Role of ABH blood group antigens in the stimulation of a DIDS-sensitive $Ca^{2+}$ influx pathway in human erythrocytes by *Ulex europaeus* agglutinin I and a monoclonal anti $A_1$ antibody (1091) 261
Engström, K.G., Sandström, P.-E. and Sehlin, J.
Volume regulation in mouse pancreatic β-cells is mediated by a furosemide-sensitive mechanism (1091) 145
Evangelopoulos, A.E., see Zevgolis, V.G. (1091) 222

Faragó, A., see Farkas, G. (1091) 81
Farkas, G., Buday, L., Antoni, F. and Faragó, A.
The 38 kDa $Ca^{2+}$/membrane-binding protein of pig granulocytes needs a high $Ca^{2+}$ concentration to be phosphorylated by protein kinase C (1091) 81
Fedorov, N.A., see Sud'ina, G.F. (1091) 257
Fehmann, H.-C., Göke, R., Göke, B., Bächle, R., Wagner, B. and Arnold, R.
Priming effect of glucagon-like peptide-1 (7-36) amide, glucose-dependent insulinotropic polypeptide and cholecystokinin-8 at the isolated perfused rat pancreas (1091) 356
Feinstein, M.B., see Banga, H.S. (1091) 115
Ferber, E., see Zor, U. (1091) 385
Fischer, G., see Zor, U. (1091) 385
Fischer, S., see Brunati, A.M. (1091) 123
Florin, I. and Thelestam, M.
ADP-ribosylation in *Clostridium difficile* toxin-treated cells is not related to cytopathogenicity of toxin B (1091) 51

Forsberg, N., see Gustafson, S. (1091) 36
Freeman, B.A., see Panus, P.C. (1091) 303
French, P.J., see Baxter, M.A. (1091) 158
Frezard, F. and Garnier-Suillerot, A.
Determination of the osmotic active drug concentration in the cytoplasm of anthracycline-resistant and -sensitive $K_{562}$ cells (1091) 29
Fuchs, K., see Deslauriers, C. (1091) 165
Fujisawa, H., see Katoh, T. (1091) 205

Gagliardino, E.E., see Gagliardino, J.J. (1091) 370
Gagliardino, J.J., Krinks, M.H. and Gagliardino, E.E.
Identification of the calmodulin-regulated protein phosphatase, calcineurin, in rat pancreatic islets (1091) 370
Gaillard, D., see Catalioto, R.-M. (1091) 364
Garner, C.M., Mills, C.O., Elias, E. and Neuberger, J.M.
The effect of bile salts on human vascular endothelial cells (1091) 41
Garnier-Suillerot, A., see Frezard, F. (1091) 29
Gibson, G.E., see Huang, H.-M. (1091) 409
Göke, B., see Fehmann, H.-C. (1091) 356
Göke, R., see Fehmann, H.-C. (1091) 356
Gomis, R., see Blachier, F. (1091) 151
Gonias, S.L., see LaMarre, J. (1091) 197
Grandin, N. and Charbonneau, M.
Changes in intracellular free calcium activity in *Xenopus* eggs following imposed intracellular pH changes using weak acids and weak bases (1091) 242
Grishin, A.V., see Nupenko, E.V. (1091) 213
Gualberto, A., see Martín, F. (1091) 110
Gustafson, S. and Forsberg, N.
Hyaluronan-binding proteins on cultured J 774 macrophages (1091) 36

Halenda, S.P., see Banga, H.S. (1091) 115
Harell, T., see Zor, U. (1091) 385
Hayes, M.A., see LaMarre, J. (1091) 197
Hazenberg, B.P.C., see Yap, S.H. (1091) 405
Heldman, E., Levine, M., Morita, K. and Pollard, H.B.
Osmotic strength differentiates between two types of calcium transport pathways regulating catecholamine secretion from cultured bovine chromaffin cells (1091) 417
Her, E., see Zor, U. (1091) 385
Herbert, J.-M. and Maffrand, J.-P.
Effect of pentosan polysulphate, standard heparin and related compounds on protein kinase C activity (1091) 432
Howell, S.L., see Persaud, S.J. (1091) 120
Huang, H.-M., Toral-Barza, L. and Gibson, G.E.
Interactions between inositol phosphates and cytosolic free calcium following bradykinin stimulation in cultured human skin fibroblasts (1091) 409
Huber, J.L., see Huber, S.C. (1091) 393
Huber, S.C. and Huber, J.L.
In vitro phosphorylation and inactivation of spinach leaf sucrose-phosphate synthase by an endogenous protein kinase (1091) 393

Iizasa, T., see Carson, D.A. (1091) 22
Ikekawa, N., see Sato, F. (1091) 188
Ishibashi, S., see Yamashita, A. (1091) 46

Jacob, J.
Fluorescence response in chlortetracycline-loaded neutrophils measures release of $Ca^{2+}$ from intracellular membrane enclosed storage sites (1091) 317
Jans, A.W.H., see Von Recklinghausen, I.R. (1091) 179
Jones, P.M., see Persaud, S.J. (1091) 120

Kaneki, M., see Sato, F. (1091) 188
Katoh, T. and Fujisawa, H.
Calmodulin-dependent protein kinase II. Kinetic studies on the interaction with substrates and calmodulin (1091) 205
Kawai, C., see Kume, N. (1091) 63
Kempson, S.A., Kunkler, K.J. and Murer, H.
Iodoacetate action on endocytic uptake of different fluid-phase markers by OK renal epithelial cells (1091) 324
Kim, H.W., see Szymańska, G. (1091) 127
Kita, T., see Kume, N. (1091) 63
Koschel, K., see Tas, P.W.L. (1091) 401
Koshkin, A.A., see Sud'ina, G.F. (1091) 257
Kranias, E.G., see Szymańska, G. (1091) 127
Krauthamer, V., see Bowmaster, T.A. (1091) 9
Krinks, M.H., see Gagliardino, J.J. (1091) 370
Kristensen, S.R.
Cell damage caused by ATP depletion is reduced by magnesium and nickel in human fibroblasts – a non-specific calcium antagonism? (1091) 285
Kume, N., Arai, H., Kawai, C. and Kita, T.
Receptors for modified low-density lipoproteins on human endothelial cells: Different recognition for acetylated low-density lipoprotein and oxidized low-density lipoprotein (1091) 63
Kunkler, K.J., see Kempson, S.A. (1091) 324
Kurokawa, T., see Yamashita, A. (1091) 46

LaMarre, J., Wollenberg, G.K., Gonias, S.L. and Hayes, M.A.
Reaction of $\alpha_2$-macroglobulin with plasmin increases binding of transforming growth factors-$\beta$1 and $\beta$2 (1091) 197
Laudanna, C., see Berton, G. (1091) 101
Laurent, G.J., see McAnulty, R.J. (1091) 231
Leader, D.P., Deana, A.D., Marchiori, F., Purves, F.C. and Pinna, L.A.
Further definition of the substrate specificity of the alpha-herpesvirus protein kinase and comparison with protein kinases A and C (1091) 426
Levine, M., see Heldman, E. (1091) 417
Limburg, P.C., see Yap, S.H. (1091) 405
Lord, J.M., see Baxter, M.A. (1091) 158
Lynch, A.M., Trebilcock, R., Tomlinson, S. and Best, L.
Studies of the mechanism of activation of HIT-T15 cells by lactate (1091) 141

Maclouf, J., see Catalioto, R.-M. (1091) 364
Maffrand, J.-P., see Herbert, J.-M. (1091) 432
Mahmood, A., see Eliakim, R. (1091) 1
Malaisse, W.J., see Blachier, F. (1091) 151
Marchiori, F., see Brunati, A.M. (1091) 123
Marchiori, F., see Leader, D.P. (1091) 426
Martín, F., Gualberto, A., Sobrino, F. and Pintado, E.
Thimerosal induces calcium mobilization, fructose 2,6-bisphosphate synthesis and cytoplasmic alkalinization in rat thymus lymphocytes (1091) 110
McAnulty, R.J., Campa, J.S., Cambrey, A.D. and Laurent, G.J.
The effect of transforming growth factor $\beta$ on rates of procollagen synthesis and degradation in vitro (1091) 231
Menegazzi, M., see Berton, G. (1091) 101
Michell, R.H., see Baxter, M.A. (1091) 158
Mills, C.O., see Garner, C.M. (1091) 41
Moreno-Aurioles, V.R. and Sobrino, F.
Glucocorticoids inhibit fructose 2,6-bisphosphate synthesis in rat thymocytes. Opposite effect of cycloheximide (1091) 96
Morita, K., see Heldman, E. (1091) 417
Moshage, H.J., see Yap, S.H. (1091) 405
Mourtada, A., see Blachier, F. (1091) 151
Murer, H., see Kempson, S.A. (1091) 324

Murono, E.P. and Washburn, A.L.
$\Delta^5$-3$\beta$-Hydroxysteroid dehydrogenase-isomerase activity in two distinct density Leydig cells from immature rats. Differences in responsiveness to human chorionic gonadotropin or 8-bromo-adenosine 3′,5′-monophosphate (1091) 55
Murphy, B.J., see Tuana, B.S. (1091) 236
Musters, R.J.P., Post, J.A. and Verkleij, A.J.
The isolated neonatal rat-cardiomyocyte used in an in vitro model for 'ischemia'. I. A morphological study (1091) 270

Nakamura, T., see Sato, F. (1091) 188
Nanes, M.S., see Rubin, J.E. (1091) 87
Naor, Z., see Zor, U. (1091) 385
Negrel, R., see Catalioto, R.-M. (1091) 364
Neuberger, J.M., see Garner, C.M. (1091) 41
Nishikawa, Y.
An FM3A mutant, G258, with a mutation that affects both cell growth and oligosaccharide-lipid synthesis (1091) 135
Nupenko, E.V., Panchenko, M.P., Starikova, M.G., Grishin, A.V. and Tkachuk, V.A.
Apparent activation of rabbit lung membrane adenylate cyclase by cytosolic proteins possessing adenylate kinase activity (1091) 213

Okamoto, Y., see Sato, F. (1091) 188
Orimo, H., see Sato, F. (1091) 188
Ouchi, Y., see Sato, F. (1091) 188

Panchenko, M.P., see Nupenko, E.V. (1091) 213
Panus, P.C., Burgess, B. and Freeman, B.A.
Characterization of cultured alveolar epithelial cell xanthine dehydrogenase/oxidase (1091) 303
Parola, M., see Albano, E. (1091) 310
Pastor-Anglada, M., see Soler, C. (1091) 193
Pauls, K.P., see Deslauriers, C. (1091) 165
Persaud, S.J., Jones, P.M. and Howell, S.L.
Activation of protein kinase C is essential for sustained insulin secretion in response to cholinergic stimulation (1091) 120
Pierce, G.N., see Tuana, B.S. (1091) 236
Pilatus, U. and Techel, D.
$^{31}$P-NMR-studies on intracellular pH and metabolite concentrations in relation to the circadian rhythm, temperature and nutrition in *Neurospora crassa* (1091) 349
Pinna, L.A., see Brunati, A.M. (1091) 123
Pinna, L.A., see Leader, D.P. (1091) 426
Pintado, E., see Martín, F. (1091) 110
Pollard, H.B., see Heldman, E. (1091) 417
Post, J.A., see Musters, R.J.P. (1091) 270
Poveda, B., see Soler, C. (1091) 193
Powell, A.D., see Deslauriers, C. (1091) 165
Purves, F.C., see Leader, D.P. (1091) 426

Raïph, S., see Brunati, A.M. (1091) 123
Rapuano, B.E. and Bockman, R.S.
Tumor necrosis factor-$\alpha$ stimulates phosphatidylinositol breakdown by phospholipase C to coordinately increase the levels of diacylglycerol, free arachidonic acid and prostaglandins in an osteoblast (MC3T3-E1) cell line (1091) 374
Reiss, N., see Zor, U. (1091) 385
Richardson, D.R. and Baker, E.
The release of iron and transferrin from the human melanoma cell (1091) 294
Roelofs, H.M.J., see Yap, S.H. (1091) 405
Rubin, J.E., Titus, L. and Nanes, M.S.
Prostaglandin $E_2$ induction of monoblastic differentiation utilizes both cAMP and non-cAMP dependent signalling systems (1091) 87

Sandström, P.-E., see Engström, K.G. (1091) 145
Sato, E., see Yamashita, A. (1091) 46
Sato, F., Okamoto, Y., Ouchi, Y., Kaneki, M., Nakamura, T., Ikekawa, N. and Orimo, H.
Biological activity of 1$\alpha$,25-dihydroxyvitamin D derivatives – 24-*epi*-1$\alpha$,25-dihydroxyvitamin D–2 and 1$\alpha$,25-dihydroxyvitamin D–7 (1091) 188
Schumacher, U., see Engelmann, B. (1091) 261
Scott, D.M., see Von Recklinghausen, I.R. (1091) 179
Sehlin, J., see Engström, K.G. (1091) 145
Sener, A., see Blachier, F. (1091) 151
Shukla, S.D., see Dhar, A. (1091) 15
Sobrino, F., see Martín, F. (1091) 110
Sobrino, F., see Moreno-Aurioles, V.R. (1091) 96
Soler, C., Poveda, B., Pastor-Anglada, M. and Soley, M.
Effect of epidermal growth factor (EGF) on gluconeogenesis in isolated rat hepatocytes. Dependency on the red-ox state of the substrate (1091) 193
Soley, M., see Soler, C. (1091) 193
Sorio, C., see Berton, G. (1091) 101
Sotiroudis, T.G., see Zevgolis, V.G. (1091) 222
Starikova, M.G., see Nupenko, E.V. (1091) 213
Sud'ina, G.F., Tatarintsev, A.V., Koshkin, A.A., Zaitsev, S.V., Fedorov, N.A. and Varfolomeev, S.D.
The role of adhesive interactions and extracellular matrix fibronectin from human polymorphonuclear leukocytes in the respiratory burst (1091) 257
Summers, B., Al-Hassan, J.M., Thomson, M., Chun, A. and Criddle, R.S.
Platelet activating factors (AGEPC) from epidermal secretions of the Arabian Gulf catfish, *Arius bilineatus*, which stimulate wound healing (1091) 329
Sunahara, R.K., see Tuana, B.S. (1091) 236
Suzuki, H., see Berton, G. (1091) 101
Szymańska, G., Kim, H.W. and Kranias, E.G.
Reconstitution of the skeletal sarcoplasmic reticulum $Ca^{2+}$-pump: influence of negatively charged phospholipids (1091) 127

Takemura, T., see Takikawa, H. (1091) 173
Takikawa, H., Tomita, J., Takemura, T. and Yamanaka, M.
Cytotoxic effect and uptake mechanism by isolated rat hepatocytes of lithocholate and its glucuronide and sulfate (1091) 173
Tas, P.W.L. and Koschel, K.
Volatile anesthetics stimulate the phorbol ester evoked neurotransmitter release from PC12 cells through an increase of the cytoplasmic $Ca^{2+}$ ion concentration (1091) 401
Tatarintsev, A.V., see Sud'ina, G.F. (1091) 257
Techel, D., see Pilatus, U. (1091) 349
Thelestam, M., see Florin, I. (1091) 51
Thomson, M., see Summers, B. (1091) 329
Titus, L., see Rubin, J.E. (1091) 87
Tkachuk, V.A., see Nupenko, E.V. (1091) 213
Tomita, J., see Takikawa, H. (1091) 173
Tomlinson, S., see Lynch, A.M. (1091) 141
Toral-Barza, L., see Huang, H.-M. (1091) 409
Trebilcock, R., see Lynch, A.M. (1091) 141
Tsunoda, Y., Williams, J.A. and DelValle, J.
Secretagogue-induced $Ca^{2+}$ oscillations in isolated canine gastric chief cells (1091) 251
Tuana, B.S., Sunahara, R.K., Murphy, B.J. and Pierce, G.N.
Two major antigens of heart sarcolemma are $Ca^{2+}$-binding glycoproteins that copurify with the dihydropyridine receptor (1091) 236
Tuijl, M.J.M., Van Bergen en Henegouwen, P.M.P., Van Wijk, R. and Verkleij, A.J.

The isolated neonatal rat-cardiomyocyte used in an in vitro model for 'ischemia'. II. Induction of the 68 kDa heat shock protein (1091) 278

Van Bergen en Henegouwen, P.M.P., see Tuijl, M.J.M. (1091) 278
Van Rijswijk, M.H., see Yap, S.H. (1091) 405
Van Wijk, R., see Tuijl, M.J.M. (1091) 278
Varfolomeev, S.D., see Sud'ina, G.F. (1091) 257
Verkleij, A.J., see Musters, R.J.P. (1091) 270
Verkleij, A.J., see Tuijl, M.J.M. (1091) 278
Von Recklinghausen, I.R., Scott, D.M. and Jans, A.W.H.
An NMR spectroscopic characterization of a new epithelial cell line, TALH-SVE, with properties of the renal medullary thick ascending limb of Henle's loop (1091) 179

Wagner, B., see Fehmann, H.-C. (1091) 356
Washburn, A.L., see Murono, E.P. (1091) 55
Wasson, D.B., see Carson, D.A. (1091) 22
Williams, J.A., see Tsunoda, Y. (1091) 251
Wollenberg, G.K., see LaMarre, J. (1091) 197

Yamanaka, M., see Takikawa, H. (1091) 173
Yamashita, A., Sato, E., Yasuda, H., Kurokawa, T. and Ishibashi, S.
Reduction of mono(ADP-ribosyl)ation of 20 kDa protein with maturation in rat testis: Involvement of guanine nucleotides (1091) 46
Yap, S.H., Moshage, H.J., Hazenberg, B.P.C., Roelofs, H.M.J., Bijzet, J., Limburg, P.C., Aarden, L.A. and Van Rijswijk, M.H.
Tumor necrosis factor (TNF) inhibits interleukin (IL)-1 and/or IL-6 stimulated synthesis of C-reactive protein (CRP) and serum amyloid A (SAA) in primary cultures of human hepatocytes (1091) 405
Yasuda, H., see Yamashita, A. (1091) 46

Zaitsev, S.V., see Sud'ina, G.F. (1091) 257
Zevgolis, V.G., Sotiroudis, T.G. and Evangelopoulos, A.E.
Phosphorylase kinase from bovine stomach smooth muscle: a $Ca^{2+}$-dependent protein kinase associated with an actin-like molecule (1091) 222
Ziegler, K., see Bertrams, A. (1091) 337
Zor, U., Her, E., Harell, T., Fischer, G., Naor, Z., Braquet, P., Ferber, E. and Reiss, N.
Arachidonic acid release by basophilic leukemia cells and macrophages stimulated by $Ca^{2+}$ ionophores, antigen and diacylglycerol: essential role for protein kinase C and prevention by glucocorticosteroids (1091) 385